

## Evaluating a Function From a Table Charge for Delivery

| Miles Traveled | Charge  |
|----------------|---------|
| $x$            | $c(x)$  |
| 1              | \$14.00 |
| 2              | \$21.00 |
| 3              | \$28.00 |
| 4              | \$35.00 |
| 5              | \$42.00 |
| 6              | \$49.00 |

## Evaluating a Function From a Table Charge for Delivery

| Miles Traveled | Charge  |
|----------------|---------|
| $x$            | $c(x)$  |
| 1              | \$14.00 |
| 2              | \$21.00 |
| 3              | \$28.00 |
| 4              | \$35.00 |
| 5              | \$42.00 |
| 6              | \$49.00 |

| Evaluate                          | Meaning  |
|-----------------------------------|--|
| $c(3) = \underline{\hspace{2cm}}$ |  |
| $c(1) = \underline{\hspace{2cm}}$ |  |
|                                   | The delivery charge for 5 miles is \$42.00.                          |
| $c(4) = \underline{\hspace{2cm}}$ |  |
|                                   | The delivery charge for $\underline{\hspace{1cm}}$ miles is \$21.00. |

| Evaluate                          | Meaning  |
|-----------------------------------|--|
| $c(3) = \underline{\hspace{2cm}}$ |  |
| $c(1) = \underline{\hspace{2cm}}$ |  |
|                                   | The delivery charge for 5 miles is \$42.00.                          |
| $c(4) = \underline{\hspace{2cm}}$ |  |
|                                   | The delivery charge for $\underline{\hspace{1cm}}$ miles is \$21.00. |